

Recent direct measurement of the Top quark mass and quasi-infrared fixed point

Biswajoy Brahmachari

Department of Physics and Astronomy
University of Maryland, College Park
MD-20740, USA.

Abstract

We note that the recent direct measurement of the top quark mass at 173.3 ± 5.6 (stat) ± 6.2 (syst) by DØ collaboration severely constrains the theoretically attractive infra-red fixed point scenario of the top quark Yukawa coupling in supersymmetric GUTs. For one-step unified models the above mentioned measurement bounds the arbitrary but experimentally determinable parameter $\tan \beta$ to the range $1.3 \leq \tan \beta \leq 2.1$. Further crunch on the top quark mass may determine $\tan \beta$ even more accurately within the fixed point scenario. On the other hand an experimental value of $\tan \beta > 2.1$ will rule out the fixed point scenario bounding $h_t^2(M_X)/4\pi$ to 0.022 from above.

There is a lot of interest among physicists in the possible measurement of the top quark mass m_t by the DØ group in the vicinity of 173 GeV [1, 2]. What does this kind of a value mean for various theoretical models trying to generate m_t ? In a theoretically well-motivated scenario [3] the top quark Yukawa coupling, and hence m_t , may get fixed at an infrared stable fixed point by the structure of the renormalization group equations (RGE) which are a coupled set of differential equations for the Yukawa couplings h_i , where $i = t, b, \tau$ with the gauge couplings. Using RGE if we evolve of the top quark Yukawa coupling h_t from a large mass scale ($M_X \simeq 10^{16} GeV$) to the scale m_t we obtain a universal value of $h_t(m_t)$ for a large domain of values of $h_t(M_X)$. The result is noticeable as it shows how the details of the possibly complicated symmetry breaking mechanisms at M_X might be obliterated by the renormalization group equations, whose fixed point structure emerges dominant at low energies. The insensitivity to the ultraviolet behaviour is a hallmark of infrared stable fixed points in all branches of physics. In this communication we want to test this attractive scenario in the light of recent direct measurement of the top quark mass by the DØ collaboration, who have reported,

$$m_t = 173.3 \pm 5.6 \text{ (stat)} \pm 6.2 \text{ (syst)} \text{ GeV.} \quad (1)$$

The prediction of the top quark mass in supersymmetric models depends on an arbitrary parameter $\tan \beta \equiv \frac{\langle H_2 \rangle}{\langle H_1 \rangle}$. Where H_2 and H_1 are the two Higgs doublets coupling to the top-quark sector and the bottom quark sectors, in contrast with the non-supersymmetric case where just one-Higgs doublet suffice, prompted by the requirement that the supersymmetric Yukawa superpotential has to be an analytic functional [4] of the superfields. In one-step

supergravity GUT models, the couplings h_t , h_b and h_τ are free parameters at the GUT scale, which are to be fixed by the low energy Fermion masses. Considering solely the third generation Yukawa couplings, at low energy the top quark mass satisfies the following relation [5] with the top quark Yukawa coupling, $\tan\beta$ and the QCD coupling α_s ;

$$m_t^{pole} = h_t(m_t) \frac{174}{\sqrt{1 + \tan^2\beta}} \left(1 + \frac{4}{3\pi} \alpha_s + 11.4 \frac{\alpha_s^2}{\pi^2}\right) \text{ GeV}. \quad (2)$$

All entries of this relation except $h_t(m_t)$ are experimentally determinate, in principle, eliciting a value of $h_t(m_t)$, related via RGE to $h_t(M_X)$. On the other hand if $h_t(M_X)$ is fixed by theoretical motivations, Eqn. (2) can be used [6] to determine $\tan\beta$, the yet un-known ‘typical’ parameter of supersymmetry. The purpose of this brief communication is to evaluate the bound on $\tan\beta$ vis-a-vis the latest measurement of the top quark mass and to identify its consequences for the fixed point scenario.

We use the two-loop RGE [7] to evolve the Yukawa as well as the gauge couplings. This determines for us the value [8] of $\alpha_s = 0.125$. Now the bounds on the top quark mass gives the upper bounds on $\tan\beta$ which have been plotted in Figure (1). The lower bound on $\tan\beta$ comes from the perturbative unitarity arguments and it depends on the top quark mass which are the shaded regions in the left of Figure (1). The lower bound $\tan\beta > 1.3$ is valid for the lower bound on the top quark mass $m_t > 161$ GeV. For the fixed point scenario $h_t^2(M_X)/4\pi \simeq 1$, we get the upper bound $\tan\beta < 2.1$. We see that apart from being attractive from a theoretical point of view, the infra-red fixed point scenario makes definitive predictions of the measurable parameter $\tan\beta$, and hence, can well be falsified. The parameter $\tan\beta$ can be determined from various studies of the supersymmetric Higgs




Figure 1: The fixed point scenario of the top quark Yukawa coupling $h_t^2(M_X)/4\pi = 1$ is the upper solid line. Lower curves represent lower values of the top quark Yukawa coupling at the GUT scale. The band at the left is the excluded region in the $\tan\beta$ plane from the perturbative unitarity of the top quark Yukawa coupling. We notice that the horizontal line at 219 GeV (old experimental upper bound) does not intersect with the predicted contours.

sector for example [10] from the heavy Higgs scalar production at future e^+e^- colliders or from inclusive semileptonic B-decays [11] which depends on the sign of the supersymmetric mixing parameter μ or from the decay of heavy charged Higgs ($m_{H^+} \gg m_t$) [12]. In any case if the value of $\tan\beta$ is determined to be greater than 2.1, the fixed point scenario in the minimal one-step unified models will be excluded.

For completeness, we note that the fixed point scenario the prediction of the b quark

mass [13] is in the range $4.29 \text{ (GeV)} < m_b(m_b) < 4.61 \text{ (GeV)}$, whereas, in the case of $h_t^2/4\pi = 0.022$ the prediction turns out to $5.44 \text{ (GeV)} < m_b(m_b) < 6.24 \text{ (GeV)}$ for all possible variation of the parameter $\tan\beta$, where the higher end of the b-quark mass range corresponds to a lower end of $\tan\beta$ domain.

Now we wish to make a remark on this deceptively simple analysis. What is new? The upper-bound on $\tan\beta$ comes from the intersection point of the experimental value of the top-quark mass (dashed lines) with the predictions. We notice that the previous [2] experimentally allowed range (dotted lines) of the top quark mass was from 158 GeV all the way to 219 GeV [CDF 176 ± 18 GeV and DØ 199 ± 20 GeV] the upper experimental bounds did not intersect the theoretical predictions. Where as, the recent direct measurements intersect the contours of theoretical predictions giving upper bounds on $\tan\beta$ corresponding to the point of intersection, which may be worth stressing, as we await more accurate determination of the top quark mass.

In conclusion, we have re-evaluated the upper bounds on $\tan\beta$ in the context of the recent direct measurement of the top-quark mass by the DØ collaboration. We have shown that if the infra red fixed point scenario is correct, $\tan\beta$ should be less than 2.1. On the contrary, if $\tan\beta > 2.1$ the fixed point scenario will be ruled out bounding the coupling $\frac{h_t^2(M_X)}{4\pi} < 0.022$; another motivation for the experimentalists to fix the value of $\tan\beta$ in supersymmetric theories.

This work has been supported by a grant from National Science Foundation. I would like

to acknowledge discussions with Prof. R. N. Mohapatra.

References

- [1] DØ collaborations, S. Abachi *et al*, DØ collaborations, hep-ex/9703008.
- [2] CDF collaborations, F. Abe *et al*, Phys. Rev. Lett. **74**, 2626 (1995); DØ collaborations, S. Abachi *et al*, Phys. Rev. Lett. **74**, 2632 (1995).
- [3] B. Pendleton and G. G. Ross, Phys. Lett. **B98**, 291, (1981); C. T. Hill, Phys. Rev. **D24**, 691, (1981); see also, M. Carena, C. E. M. Wagner, Nucl. Phys. **B452**, 45 (1995).
- [4] For a review see, P. West, “*Introduction to supersymmetry and supergravity*” (World Scientific, Singapore, 1990)
- [5] R. Tarrach, Nucl. Phys. **B183**, 384 (1981); H. Arason, D. J. Castano, B. Keszthelyi, S. Mikaelian, E.J. Piard, P. Ramond, B.D. Wright, Phys. Rev. **D46**, 3945 (1992).
- [6] See reference [9] and also M. Carena, S. Dimopoulos, C. E. M. Wagner, S. Raby, Phys. Rev. **D52**, 4133 (1995).
- [7] By D. R. T. Jones Phys.Rev. **D25**, 581, (1982); M. B. Einhorn, D. R. T. Jones, Nucl. Phys. **B196**, 475 (1982). Expressions of anomalous dimensions are given in, M. Bastero-Gil and B. Brahmachari Nucl. Phys. **B482**, 39 (1996).

- [8] For detailed evaluation of α_s in supersymmetric GUTs see for example, J. Bagger, K. Matchev and D. Pierce, Phys. Lett. **B 348**, 443 (1995); P. H. Chankowski, Z. Pluciennik and S. Pokorski, Nucl. Phys. **B 439**, 23 (1995); M. Bastero-Gil and J. Pérez-Mercader, Nucl Phys. **B450**, 21 (1995).
- [9] See for example: V. Barger, M. S. Berger and P. Ohmann, Phys. Rev. **D 47**, 1093, (1993). For a brief summary of the RGE results in SUSY GUTs see, V. Barger, M. S. Berger, P. Ohmann and R. N. J. Phillips, University of Wisconsin Madison Report No. MAD/PH/803 (unpublished) and references therein; Phys. Lett. **B314**, 351, (1993).
- [10] Jonathan. L. Feng, Takeo Moroi, hep-ph/9612333.
- [11] J.A. Coarasa, R. A. Jimenez, J. Sola, hep-ph/9701392.
- [12] A. Czarnecki, J. L. Pinfold, Phys. Lett. **B328**, 427 (1994).
- [13] J.E. Bjorkman, D.R.T. Jones; Nucl. Phys. **B259**, 533 (1985); B. Ananthanarayan, G. Lazarides and Q. Shafi, Phys. Rev. **D 44**, 1613 (1990); P. Langacker, N. Polonsky, Phys. Rev. **D49**, 1454 (1994). R. Rattazzi, U. Sarid, L. J. Hall, SU-ITP-94-15, Presented at 2nd IFT Workshop on Yukawa Couplings and the Origins of Mass, Feb. 11-13, 1994, Gainesville, Florida. Also, Arason *et al*, in Ref. [5]. Nir Polonsky, Phys. Rev. **D54**, 4537 (1996).